

Osmosensitivity Associated with Insertions in *argP* (*iciA*) or *glnE* in Glutamate Synthase-Deficient Mutants of *Escherichia coli*

Madhusudan R. Nandineni,^{1,2} Rakesh S. Laishram,² and J. Gowrishankar^{2*}

Centre for Cellular and Molecular Biology,¹ and Laboratory of Bacterial Genetics, Centre for DNA Fingerprinting and Diagnostics,² Hyderabad, India

Received 1 April 2004/Accepted 7 July 2004

An ampicillin enrichment strategy following transposon insertion mutagenesis was employed to obtain NaCl-sensitive mutants of a *gltBD* (glutamate synthase [GOGAT]-deficient) strain of *Escherichia coli*. It was reasoned that the *gltBD* mutation would sensitize the parental strain even to small perturbations affecting osmotolerance. Insertions conferring an osmosensitive phenotype were identified in the *proU*, *argP* (formerly *iciA*), and *glnE* genes encoding a glycine betaine/proline transporter, a LysR-type transcriptional regulator, and the adenylyltransferase for glutamine synthetase, respectively. The *gltBD*⁺ derivatives of the strains were not osmosensitive. The *argP* mutation, but not the *glnE* mutation, was associated with reduced glutamate dehydrogenase activity and a concomitant NH₄⁺ assimilation defect in the *gltBD* strain. Supplementation of the medium with lysine or a lysine-containing dipeptide phenocopied the *argP* null mutation for both osmosensitivity and NH₄⁺ assimilation deficiency in a *gltBD* background, and a dominant gain-of-function mutation in *argP* was associated with suppression of these lysine inhibitory effects. Osmosensitivity in the *gltBD* strains, elicited either by lysine supplementation or by introduction of the *argP* or *glnE* mutations (but not *proU* mutations), was also correlated with a reduction in cytoplasmic glutamate pools in cultures grown at elevated osmolarity. We propose that an inability to accumulate intracellular glutamate at high osmolarity underlies the osmosensitive phenotype of both the *argP* *gltBD* and *glnE* *gltBD* mutants, the former because of a reduction in the capacity for NH₄⁺ assimilation into glutamate and the latter because of increased channeling of glutamate into glutamine.

When bacteria are exposed to environments with elevated osmolarity, there is a passive outflow of water from the intracellular compartment, which results in a growth-inhibiting loss of cell turgor and a reduction in the cytoplasmic volume. The set of metabolic changes that occur under these conditions to restore bacterial growth is referred to as osmoregulation. In enterobacteria such as *Escherichia coli* and *Salmonella enterica* serovar Typhimurium, the osmoregulatory processes that have been identified so far include cell envelope alterations and the cytoplasmic accumulation (to concentrations that approach several hundred millimolar) of K⁺ ions, glutamate, trehalose, proline, and glycine betaine (reviewed in references 12 and 13). Of these, the accumulation of glutamate and trehalose is mediated by increased synthesis of these compounds, whereas accumulation of K⁺, proline, and glycine betaine is achieved by increased uptake from the culture medium. Glutamate serves as a cytoplasmic counterion for K⁺ in osmotically stressed cells, and the available evidence suggests that perturbations in accumulation of either one of these molecules adversely affects the accumulation of the other (31, 43). Proline and glycine betaine are also referred to as osmoprotectants, because at low concentrations they dramatically enhance the growth rates of enterobacteria in media with elevated osmolarity.

In addition to its perceived role in osmoregulation, glutamate is also a central player in global nitrogen metabolism

(reviewed in references 35 to 37); 75 to 90% of all cellular nitrogen is assimilated via glutamate. NH₄⁺ is the preferred nitrogen source for *E. coli*, and it is assimilated into glutamate through two pathways. One of these pathways is the glutamate dehydrogenase (GDH) pathway, in which 2-oxoglutarate undergoes reductive condensation with NH₄⁺, yielding glutamate. The second pathway is the two-step glutamine synthetase (GS)-glutamate synthase (GOGAT) pathway, in which glutamine that is synthesized in the first step from one molecule of NH₄⁺ and glutamate (in the presence of ATP) is involved in a reductive reaction with 2-oxoglutarate, which yields two molecules of glutamate. GDH, GS, and GOGAT are encoded by the *gdhA*, *glnA*, and *gltBD* genes, respectively.

The GDH pathway is functional for nitrogen assimilation in media containing ≥ 1 mM NH₄⁺. In media with limiting NH₄⁺ concentrations or with alternative poor nitrogen sources, nitrogen assimilation into glutamate is mediated through the GS-GOGAT pathway, whose regulation in turn is tied to the complex cascade of nitrogen regulation that is referred to as Ntr (35–37). In NH₄⁺-replete media, *glnA* is expressed only at basal levels; in addition, there is a reduction in the catalytic activity of GS resulting from the adenylylation of its homopolymeric subunits by the *glnE*-encoded adenylyltransferase. The residual activity of GS is then sufficient to meet the cell's anabolic requirement for glutamine for protein synthesis. On the other hand, in low-NH₄⁺ medium or during growth on poor nitrogen sources, expression of the Ntr regulon (of which *glnA* is a member) is activated, and in addition GS is deadenylylated by GlnE; the vastly increased activity of GS is now able

* Corresponding author. Mailing address: Laboratory of Bacterial Genetics, Centre for DNA Fingerprinting and Diagnostics, ECIL Road, Nacharam, Hyderabad 500 076, India. Phone: 91-40-27155609. Fax: 91-40-27155610. E-mail: shankar@cdf.org.in.

to catalyze sufficient glutamine synthesis to meet the cell's nitrogen assimilatory requirement.

With high exogenous levels of NH_4^+ , glutamate accumulation at high osmolarity is not affected by single mutations that block either biosynthetic pathway (3, 31), but in low- NH_4^+ media, GOGAT-deficient mutants are osmosensitive (14, 43). A correlation between osmotolerance and NH_4^+ assimilation efficiency in GOGAT-deficient strains was established in a previous study (39). The data obtained previously provided indirect genetic evidence that increased glutamate synthesis is necessary for optimal growth under hyperosmotic stress conditions.

In the present study, we performed transposon insertion mutagenesis of a GOGAT-deficient (*gltBD*) strain to identify mutants that are osmosensitive on NH_4^+ -replete medium. The rationale was that the *gltBD* strain would be sensitized to even small perturbations that affect osmotolerance (given that multiple additive mechanisms operate for osmoregulation). Furthermore, it was reasoned that transposon insertions would generate null mutations and that the possibility of obtaining conditional-lethal salt-sensitive mutants with mutations in essential housekeeping genes would be avoided (12).

We report here identification of insertions in *glnE*, as well as in *argP*, each of which acts synergistically with *gltBD* in conferring an osmosensitive phenotype. As mentioned above, *glnE* is involved in covalently modifying and thus modulating GS activity. The *argP* gene (also called *iciA*) encodes a LysR-type transcriptional regulator that has previously been implicated in the regulation of arginine transport (6–8, 34) and in the control of chromosomal DNA replication initiation (20, 21, 42) in *E. coli*.

MATERIALS AND METHODS

Bacterial strains and growth media. Genotypes of *E. coli* K-12 strains are shown in Table 1. The routine rich and minimal growth media used were Luria-Bertani medium (33) and 0.2% glucose–minimal A medium (33), respectively, and the incubation temperature was 37°C. Minimal A medium contains 15 mM NH_4^+ and hence is considered to be NH_4^+ replete. Growth in liquid cultures was monitored with a Klett-Summerson photoelectric colorimeter. NH_4^+ assimilation growth experiments were performed by using W salts basal medium (40), to which 0.4% glucose was added as a carbon source and the desired concentration of ammonium sulfate was added as a nitrogen source. The medium was supplemented with lysine (Lys), lysylalanine (Lys-Ala), histidylalanine, or glycine betaine (each at a concentration of 1 mM) and with aspartate (0.2%). Unless otherwise indicated, the concentrations of antibiotics used were the concentrations described previously (39).

Phages and plasmids. The transposon phage $\lambda\text{placMu55(Kan)}$ encoding kanamycin resistance and its helper phage λpMu507 have been described previously (29), as have the plasmid vectors (i) pCL1920 (pSC101 based, low copy number, spectinomycin and streptomycin resistant) (27), (ii) pACYC184 (p15A based, medium copy number, chloramphenicol and tetracycline resistant) (9), (iii) pBR329 (pMB9 based, high copy number, ampicillin, tetracycline, and chloramphenicol resistant) (10), and (iv) pBluescriptII-KS (pMB9 based, very high copy number, ampicillin resistant) (Stratagene, La Jolla, Calif.).

Plasmid pHYD909 was constructed by subcloning a 1.6-kb BamHI-HindIII fragment encoding an N-terminally truncated FNR protein from plasmid pGS198 (41) into the appropriate sites of pACYC184. Plasmids pHYD916, pHYD942, and pHYD943 are derivatives of pCL1920 carrying the following subcloned fragments from recombinant λ phage clones of the ordered genomic library of Kohara et al. (23): in pHYD916, 5.2-kb BamHI-KpnI fragment with *glnE*⁺ from λ 508 in the corresponding sites of the vector; in pHYD942, 0.78-kb PstI-HpaI fragment with *apaG*⁺ from λ 105 in the BamHI-SalI sites of the vector; and in pHYD943, 1.06-kb ClaI-EcoRI fragment with *apaH*⁺ from λ 105 in the BamHI-SalI sites of the vector (the last two via an intermediate step of cloning into pBluescriptII-KS). Plasmids pHYD915, pHYD953, and pHYD954 have

TABLE 1. *E. coli* K-12 strains^a

Strain	Genotype ^b
MC4100	$\Delta(\text{argF-lac})\text{U169 rpsL150 relA1 araD139 ffbB5301 deoC1 ptsF25}$
GJ2529	MC4100 $\Delta\text{gltBDF500 zha-6::Tn10}$
GJ2530	GJ2529 <i>fnr::Ω</i>
GJ2560	GJ2529 <i>proU610::$\lambda\text{placMu55(Kan)}$</i> ^d
GJ4534	MC4100 <i>proU610::$\lambda\text{placMu55(Kan)}$</i>
GJ4536	MC4100 <i>argP202::$\lambda\text{placMu55(Kan)}$</i>
GJ4540	MC4100 <i>argP203::$\lambda\text{placMu55(Kan)}$</i>
GJ4544	MC4100 <i>proU611::$\lambda\text{placMu55(Kan)}$</i>
GJ4545	MC4100 <i>glnE463::$\lambda\text{placMu55(Kan)}$</i>
GJ4547	MC4100 <i>glnE464::$\lambda\text{placMu55(Kan)}$</i>
GJ4549	MC4100 <i>glnE465::$\lambda\text{placMu55(Kan)}$</i>
GJ4652	MC4100 $\Delta\text{gltBDF500}$
GJ4654	GJ4652 <i>argP202::$\lambda\text{placMu55(Kan)}$</i>
GJ4655	GJ4652 <i>argP203::$\lambda\text{placMu55(Kan)}$</i>
GJ4661	GJ4652 <i>apaG611::$\lambda\text{placMu55(Kan)}$</i>
GJ4663	GJ4652 <i>glnE463::$\lambda\text{placMu55(Kan)}$</i>
GJ4664	GJ4652 <i>glnE464::$\lambda\text{placMu55(Kan)}$</i>
GJ4735	GJ4652 <i>apaH::Kan</i>
GJ4842	GJ4652 $\Delta\text{ruvABC::Cm}$
GJ4843	GJ4652 <i>recA56 srl-300::Tn10</i>
GJ4891	GJ4663 <i>argP202::$\lambda\text{placMu55(Kan)}$</i> $\Delta\text{dsbC::Cm}$
GJ4928	GJ4652 <i>proU611::$\lambda\text{placMu55(Kan)}$</i>
GJ4929	GJ4652 <i>glnE465::$\lambda\text{placMu55(Kan)}$</i>

^a Strain MC4100 was obtained from our lab stock collection (17). All other strains were obtained in this study.

^b Genotype designations are those described by Berlyn (1).

^c The Ω insertion cassette encodes streptomycin and spectinomycin resistance.

^d Based on the molecular characterization data described in Table 2, the *proU610* insertion is situated in *proV*, the first of the three constituent genes of the *proU* operon (1).

been described previously (34) and carry *argP*⁺ cloned in pCL1920, the dominant gain-of-function *argP* (*argP*-S94L) allele cloned in pBR329, and the *argO cis* regulatory region cloned in pBluescriptII-KS, respectively.

Isolation and molecular genetic characterization of NaCl-sensitive mutants. By using the infection protocol described previously (29), populations of clones were obtained that carried random transpositions of the *lac* fusion phage $\lambda\text{placMu55(Kan)}$ in the genome of strain GJ2530/pHYD909. Approximately 10⁵ independent Kan^r insertions were obtained in each of two different experiments. Each of the populations was inoculated at a concentration of 10⁷ cells per ml into 10 ml of glucose–minimal A medium supplemented with 0.7 M NaCl and glycine betaine and incubated with shaking until the *A*₆₀₀ was around 0.1; after this ampicillin was added to a concentration of 100 $\mu\text{g/ml}$. Surviving cells after incubation for an additional 4 h were harvested by filtration and grown overnight in glucose–minimal A medium, and the cycle used for ampicillin enrichment for cells unable to grow in NaCl-supplemented medium was repeated. The cultures were then plated on glucose–minimal A medium, and individual colonies were tested for NaCl sensitivity.

Molecular mapping of the mutation in each of the derivatives in which the NaCl sensitivity phenotype was 100% cotransducible with the Kan^r marker of the transposon was performed by using an inverse PCR approach. Chromosomal DNA was digested to completion with HhaI, ligated at a high dilution, and then subjected to PCR amplification with a pair of divergently oriented primers (MuC1 [5'-TGCGTTTTTCTTCAGGTAATG-3'] and MuC2 [5'-TCCCGAAT AATCCAATGTCCTCCCG-3']) from the sequence at the *c* end of phage Mu. This procedure was expected to amplify the HhaI fragment at the junction of the Mu *c* end of the insertion and the chromosome. (If the terminal nucleotide pair at the *c* end of Mu is counted as bp 1, the MuC1 and MuC2 primers are designed to read inward from bp 438 and outward from bp 56, respectively, and the HhaI site is at bp 704.) Following agarose gel electrophoresis, the PCR products were eluted from the gel and sequenced with an automated DNA sequencer by using primer MuC2. The identity of the mutated gene was then established by a BLAST search analysis of the sequence determined against the *E. coli* genome sequence (2).

GDH assays. For GDH assays, cultures initially grown in glucose–minimal A medium with glycine betaine to an *A*₆₀₀ of around 0.2 were each split into two parts. To one part (high osmolarity grown) NaCl was added, from a 2.4 M stock

TABLE 2. Molecular characterization and NaCl tolerance of λ placMu55(Kan) insertion mutants

Insertion allele	Genomic position (bp) ^a	Mu orientation ^b	<i>lacZ</i> orientation ^c	Distance from start of gene (total gene length) (bp) ^d	NaCl tolerance ^e	
					<i>gltBD</i> (strain)	<i>gltBD</i> ⁺ (strain)
<i>apaG11</i>	51249	CCW	–	358 (375)	S (GJ4661)	ND ^f
<i>argP202</i>	3057872	CCW	+	104 (891)	S (GJ4654)	T(GJ4536)
<i>argP203</i>	3057754	CCW	+	–19 (891)	S (GJ4655)	T(GJ4540)
<i>glnE463</i>	— ^g	—	–	—	S (GJ4663)	T(GJ4545)
<i>glnE464</i>	3196273	CW	+	1,014 (2,838)	S (GJ4664)	T(GJ4547)
<i>glnE465</i>	3194695	CCW	–	2,588 (2,838)	S (GJ4929)	T(GJ4549)
<i>proU610</i>	2803134	CCW	+	303 (1,200)	S (GJ2560)	T(GJ4534)
<i>proU611</i>	— ^g	—	+	—	S (GJ4928)	T(GJ4544)

^a The nucleotide position in the *E. coli* genome at the junction with the Mu *c* end of each insertion is indicated. The genome sequence is from reference 2 (GenBank accession number U00096).

^b CW and CCW refer to the two alternative orientations, in which the Mu *S* end of the insertion is clockwise and counterclockwise, respectively, from the Mu *c* end in the *E. coli* chromosome.

^c + and –, orientations in which the direction of the *lacZ* reporter gene within the transposon is the same as and opposite that of the host gene disrupted by the insertion, respectively.

^d Distance from the start codon to the Mu insertion site and length of the wild-type gene. The fact that Mu transposition is associated with a 5-bp target sequence duplication (33) was also taken into account when the distance was calculated.

^e Growth of the *gltBD* and *gltBD*⁺ derivatives was compared with that of the control strains GJ4652 (*gltBD*) and MC4100 (*gltBD*⁺) on glucose-minimal A agar plates supplemented with glycine betaine and several concentrations of NaCl. S, sensitive; T, tolerant.

^f ND, not determined.

^g —, not molecularly characterized (the Mu *c* end in *glnE463* is 6 bp away from an *HhaI* site in *glnE*).

solution prepared in glucose-minimal A medium with glycine betaine, to a final concentration of 0.6 M, and incubation was continued. The other part (low osmolarity grown) was incubated without further manipulation. All cultures were harvested at an A_{600} of around 0.5.

Cell extracts for determination of GDH activity were prepared as described previously (19), with the following modifications: (i) after harvested cells were washed, they were resuspended in 4 ml of ice-cold 50 mM Tris-Cl buffer (pH 7.6)–10 mM β -mercaptoethanol and (ii) cells were disrupted by sonication. The extracts were used for enzyme activity measurements immediately. The assay method was the method described by Meers et al. (32), except that the reactions were performed at room temperature in 0.5-ml (total volume) mixtures. Protein concentrations in cell extracts were determined by the method of Bradford (4). Enzyme specific activities were expressed in milliunits per milligram of protein in the cell extracts (after correction for endogenous NADPH oxidase activity); 1 U was defined as the amount of enzyme required to oxidize 1 μ mol of NADPH (extinction coefficient at 340 nm, 6,220 M⁻¹ cm⁻¹) per min.

Estimation of intracellular glutamate and glutamine pools. For estimation of intracellular glutamate and glutamine pools, cultures grown at low and high osmolarities were prepared in the way described above for the GDH assays. (When cultures were supplemented with Lys or Lys-Ala, they were supplemented at all stages of growth.) A 0.2-ml portion of each culture was added to 0.8 ml of ice-cold methanol as recommended in the no-harvest method described by Kustu and coworkers (14, 25, 43). Cell debris in the mixture was removed by low-speed centrifugation, and the supernatant was lyophilized, resuspended in 0.05 ml of water, and filtered through a 0.45- μ m-pore-size filter. Proteins in the sample were precipitated by addition of 0.05 ml of 10% trichloroacetic acid, and the supernatant obtained after low-speed centrifugation was lyophilized. The content of glutamate and glutamine in the sample was then estimated, after precolumn derivatization with phenylisothiocyanate, by reverse-phase high-performance liquid chromatography with a Waters Pico · Tag column for free amino acids (Millipore Corp, Milford, Mass.) and detection of absorbance at 254 nm; the protocols for sample derivatization and column chromatography were similar to those described in the manufacturer's instructions. As standards for quantitation, 250 pmol of glutamate and 250 pmol of glutamine were injected after derivatization into the column, and, under the conditions employed, they eluted at 2.99 and 5.02 min, respectively.

For determination of the dry weight of bacteria in the culture, a 15- to 20-ml portion of a culture was filtered through a preweighed 0.45- μ m-pore-size filter (diameter, 47 mm); the filter was washed with an equal volume of water and then dried at room temperature to a constant weight. By this method, we calculated the mean bacterial dry weights to be 0.82 and 0.46 mg ml⁻¹ A_{600} unit⁻¹ for cultures grown in glucose-minimal A medium with glycine betaine and in glucose-minimal A medium with glycine betaine and 0.6 M NaCl, respectively.

Other methods. Mutations were transferred between strains by P1 transduction (17). Recombinant DNA procedures were performed as described previously (38).

RESULTS

Isolation of NaCl-sensitive mutants from *gltBD* strain. As described above, an ampicillin enrichment strategy was employed with a population of transposon-mutagenized clones of a *gltBD* strain in order to identify insertions that conferred an NaCl-sensitive phenotype on glycine betaine-supplemented medium. Since glycine betaine is osmoprotective for *E. coli* (12, 13), its inclusion permitted ampicillin selection to be performed at a sufficiently high NaCl concentration, 0.7 M. (The starting strain also carried a Δ *fnr* mutation, which is known to confer osmosensitivity in the *gltBD* background [39], along with a plasmid encoding an N-terminally truncated FNR protein that is proficient for complementing the *fnr* mutant for osmosensitivity; however, our subsequent studies [data not shown] indicated that neither the *fnr* mutation nor its complementing plasmid was relevant to the phenotypes described below.) The Kan^r insertions were each transduced into Δ *gltBD* strain GJ4652 in order to establish that NaCl sensitivity is 100% linked to Kan^r.

The sites of transposon insertions in the mutants were then determined by two complementary approaches, inverse PCR (as described above) and transductional mapping (Tables 2 and 3). One Kan^r insertion in *apaG*, two Kan^r insertions each in *proU* and *argP*, and three Kan^r insertions in *glnE* were identified that conferred an NaCl-sensitive phenotype in the *gltBD* background.

On the *E. coli* chromosome, the *apaG* gene is upstream of and is cotranscribed with *apaH*. The latter gene encodes an Ap₄A hydrolase (28). We were able to demonstrate by using the following two criteria that the NaCl-sensitive phenotype conferred by the *apaG* insertion in the *gltBD* strain is the result of a polar effect on *apaH* expression (data not shown): (i) an *apaH*::Kan mutation described previously (28) conferred NaCl sensitivity in the *gltBD* strain, which was more pronounced than the NaCl sensitivity observed with *apaG*; and (ii) NaCl tolerance in both the *apaG gltBD* (GJ4661) and *apaH gltBD* (GJ4735) strains was restored upon introduction of the mini-

TABLE 3. P1 transductional mapping of λ placMu55(Kan) insertions^a

λ placMu55(Kan) allele(s) (min)	Donor marker (min)	% Linkage
<i>apaG11</i> (1.1)	<i>zab-3051::Tn10</i> (1.8)	22
<i>argP202</i> , -203 (65.9)	<i>ΔdsbC::Cm</i> (65.4)	12–33
	<i>galP::Tn10</i> (66.5)	16–19
	<i>mutY::Tn10</i> (66.8)	6–10
<i>glnE463</i> , -464, -465 (68.9)	<i>metC162::Tn10</i> (67.9)	11–16
<i>proU610</i> , -611 (60.4)	<i>zga-900::Tn10</i> (60.3) ^b	67–100

^a In each cross, linkage was determined as the proportion of Kan^s derivatives among the transductants inheriting the donor marker (chloramphenicol resistance in the case of Δ dsbC::Cm and tetracycline resistance in the case of all other markers). For the loci with two or more λ placMu55(Kan) insertion alleles, the range of linkage values obtained is indicated.

^b The *zga-900::Tn10* insertion was designated *zfi-900::Tn10* in a previous study (17).

mal *apaH*⁺ plasmid pHYD943 but not upon introduction of the minimal *apaG*⁺ plasmid pHYD942. However (in a finding which is different from the finding reported by Leveque et al. [28] in a different *E. coli* strain background), we observed that the *apaG* and *apaH* insertions (in both *gltBD* and *gltBD*⁺ derivatives of MC4100) were associated with slow growth even in glucose-minimal A and succinate-minimal A media that were not supplemented with NaCl (data not shown). The *apaH* mutant also exhibited a reduced growth rate in hypotonic medium, such as 0.2 \times minimal A medium with NH₄⁺ added to a concentration of 15 mM and glucose. Therefore, we could not exclude the possibility that the NaCl sensitivity of the *apaG* or *apaH* mutants was just a manifestation of their nonspecific slow-growth phenotype, not unlike the behavior of other slowly growing control strains, such as *recA* or *ruvABC* strains (see below), which are not expected to be affected in NaCl tolerance. Accordingly, further studies with the *apaGH* mutants were not pursued.

***argP gltBD* and *glnE gltBD* mutants are osmosensitive.** By using plate growth tests, insertions in each of the other three loci (*proU*, *argP*, and *glnE*) were shown (i) to confer NaCl sensitivity only in the *gltBD* background and not when they were transduced into the isogenic *gltBD*⁺ strain MC4100 (Table 2) and (ii) not to affect growth on media not supplemented with NaCl in either strain background (data not shown). Since the role in osmoregulation of the glycine betaine/proline transporter encoded by the three constituent genes (*proV*, *proW*, and *proX*) of the *proU* operon is well established (12, 13), the two new *proU* insertion mutants identified in this study were not characterized further.

Of the two insertions in *argP*, the one located in the interval between the putative promoter and the start of the structural gene (*argP203*) conferred a less severe phenotype than the one in the coding region of the gene (*argP202*). All three *glnE gltBD* mutants were equally compromised for NaCl tolerance. The *argP gltBD* and *glnE gltBD* mutants were complemented for NaCl tolerance by pSC101-based plasmids carrying the *argP*⁺ (pHYD915) and *glnE*⁺ (pHYD916) genes, respectively. Neither of these plasmids (nor derivatives of pBR329 with the cloned *argP*⁺ or *glnE*⁺ genes) had any significant effect on NaCl tolerance of *gltBD*⁺ or *gltBD* strains that were *argP*⁺ *glnE*⁺ (data not shown).

We compared the growth rates of *argP202 gltBD* and

glnE463 gltBD mutants (test strains) on the one hand and the *gltBD*, *recA gltBD*, and Δ *ruvABC gltBD* strains (control strains) on the other hand in (i) glucose minimal medium, (ii) succinate minimal medium, and (iii) glucose minimal medium rendered hyperosmolar with either 0.8 M NaCl or 1.2 M glucose (Fig. 1). It is known that the *gltBD* mutant is only marginally osmosensitive compared with the wild-type strain (14, 39, 43). The *recA* and *ruvABC* derivatives were chosen as mutants whose growth deficiency is not related to the osmolarity of the growth medium, and likewise succinate minimal medium was chosen to represent a growth stress not related to osmotic stress. It was reasoned that a true osmosensitive mutant was a mutant that was specifically defective for growth only in media with elevated osmolarity. By using these criteria, both test strains were classified as osmosensitive, since (i) in the succinate minimal medium, the *argP gltBD* strain's growth rate was similar to that of the *gltBD* single mutant, while the *glnE gltBD* strain grew at least as well as the *recA gltBD* derivative; and (ii) in the 0.8 M NaCl- or 1.2 M glucose-supplemented cultures, the test strains exhibited significantly reduced growth rates compared with all three control strains (Fig. 1).

The test strains were also significantly more sensitive than the wild-type or *gltBD* control strains to other ionic or nonionic impermeant solutes, including KCl, K₂SO₄, NH₄Cl, (NH₄)₂SO₄, and sucrose, but they were not more sensitive than the control strains to equiosmolar concentrations of freely permeable solutes, such as glycerol or ethylene glycol (data not shown). These observations indicated that the test strains are deficient for osmotic stress adaptation. The *argP glnE gltBD* triple mutant GJ4891 was even more osmosensitive than the *argP gltBD* and *glnE gltBD* double mutants (data not shown).

Exacerbation by *argP* of NH₄⁺ assimilation deficiency in *gltBD* mutant. As mentioned above, a *gltBD* mutant (in which the GDH pathway of NH₄⁺ assimilation alone is functional) is not significantly nitrogen limited in a medium containing NH₄⁺ at a concentration of 1 mM or more (35–37). We observed that the *argP gltBD* mutant was unable to grow even on medium containing 2 mM NH₄⁺, whereas a *glnE gltBD* strain and the *gltBD* single mutant were able to grow on this medium (Fig. 2). On the other hand, with 15 mM NH₄⁺ (Fig. 2) or with aspartate as the nitrogen source (which readily donates its amino group for glutamate synthesis) (data not shown), the growth of the *argP gltBD* mutant was similar to that of the *gltBD* strain. Thus, osmosensitivity in the *argP gltBD* strain could be correlated with a deficiency in NH₄⁺ assimilation into glutamate. The growth of the *argP gltBD*⁺ strain was not affected even in media containing <1 mM NH₄⁺ (data not shown).

Reduced GDH activity in *argP gltBD* strain. GDH specific activities were then determined by the method described above for wild-type strain MC4100, as well as its *gltBD*, *argP gltBD*, and *glnE gltBD* derivatives, which were grown in NH₄⁺-replete cultures at low or high osmolarity (Table 4). We observed that osmotic stress was associated with a twofold increase in GDH specific activity in GJ4652 (*gltBD*) but not in MC4100 (wild type), a difference that may perhaps reflect the existence in the latter strain of redundant pathways for increased glutamate synthesis for osmoregulation. The GDH activities in cultures of the *glnE gltBD* strain GJ4663 at both low and high osmolarities were not significantly different from those in cultures of




FIG. 1. Osmosensitivity of *glnE gltBD* and *argP gltBD* mutants. Semilogarithmic growth curves were plotted for strains GJ4652 (*gltBD*), GJ4654 (*gltBD argP*), GJ4663 (*gltBD glnE*), GJ4842 (*gltBD ruvABC*), and GJ4843 (*gltBD recA*) in different media after 1:100 inoculation from cultures grown to the stationary phase in glucose-minimal A medium. The media employed were 0.2% glucose-minimal A medium (Glc-MA), 0.2% succinate-minimal A medium (Suc-MA), glucose-minimal A medium supplemented with glycine betaine and 0.8 M NaCl (0.8 M NaCl), and glucose-minimal A medium supplemented with glycine betaine and 1.2 M glucose (1.2 M Glc).

GJ4652. On the other hand, for the *argP gltBD* strain GJ4654, the GDH activity in the low-osmolarity culture was only one-half that of GJ4652, and furthermore there was no increase in enzyme activity in the strain following imposition of NaCl stress. These results established that NH_4^+ assimilation defi-

ciency and reduced GDH activity are correlated in the *argP gltBD* strain.

Osmosensitivity is correlated with decreased glutamate accumulation in both *argP gltBD* and *glnE gltBD* mutants. The fact that the *argP* and *glnE* mutants were affected for NH_4^+




FIG. 2. Effects of *argP* mutations and Lys or Lys-Ala supplementation on NH_4^+ assimilation in *gltBD* derivatives. Each strain was streaked on a pair of agar plates containing W salts medium supplemented with 2 mM NH_4^+ (top row) and 15 mM NH_4^+ (bottom row) (NH_4^+ was added as ammonium sulfate to concentrations of 1 and 7.5 mM, respectively). Additional genetic markers present in the *gltBD* derivatives are indicated above each pair of panels, and the strains employed were GJ4652 (Nil), GJ4654 (*argP*), GJ4663 (*glnE*), GJ4652/pBR329 (vector), and GJ4652/pHYD953 (*argP^d*, that is, with the dominant *argP*-S94L mutation). The plasmid-bearing strains were streaked on ampicillin-containing medium with Lys or Lys-Ala, as indicated. The plates were incubated for 24 h.

TABLE 4. GDH activities in *argP gltBD* and *glnE gltBD* derivatives^a

Strain	Genotype	GDH sp act (μg mg of protein ⁻¹)	
		Low osmolarity	High osmolarity
MC4100	Wild type	90	105
GJ4652	<i>gltBD</i>	111	224
GJ4654	<i>argP gltBD</i>	54	46
GJ4663	<i>glnE gltBD</i>	107	170

^a GDH specific activities were measured in cells that were grown as described in the text and were harvested from (i) glucose-minimal A medium with glycine betaine (low osmolarity) or (ii) glucose-minimal A medium with glycine betaine and 0.6 M NaCl (high osmolarity). Each value is the average of at least two independent determinations, and the variation between individual measurements was less than 20%.

assimilation and GS regulation, respectively, combined with the fact that each of the mutations acted synergistically with *gltBD* for osmosensitivity, made it likely that insufficient glutamate accumulation was responsible for the osmosensitive phenotypes of the mutants. Accordingly, we measured the intracellular glutamate and glutamine pools in the cultures of these mutants (along with those of the wild-type, *gltBD*, and *proU gltBD* strains as controls) grown in NH₄⁺-replete medium at low and elevated osmolarities (Table 5).

Consistent with previous reports (14, 30, 31, 43), there was substantial accumulation of glutamate in cells of wild-type strain MC4100 grown at the elevated osmolarity, which was accompanied by a small increase in the size of the intracellular glutamine pool. Compared to the level in MC4100, the intracellular glutamate levels in NaCl-grown cultures of the *gltBD* and *proU gltBD* mutants were about 40% lower, whereas the reduction was even more pronounced (around 80%) for the *argP gltBD* and *glnE gltBD* strains. At the same time, the size of the glutamine pool in high-osmolarity-grown cells of the *glnE*

TABLE 5. Glutamate and glutamine pools in *gltBD* derivatives^a

Strain genotype ^b	Supplement	Concn (nmol mg [dry wt] of cells ⁻¹) of:			
		Glutamate		Glutamine	
		Low osmolarity	High osmolarity	Low osmolarity	High osmolarity
Wild type	None	36	212	5	26
<i>gltBD</i>	None	26	117	7	43
<i>proU gltBD</i>	None	30	118	5	36
<i>argP gltBD</i>	None	18	50	7	40
<i>glnE gltBD</i>	None	21	39	19	236
<i>gltBD</i>	Lys	ND ^c	46	ND	21
<i>gltBD</i>	Lys-Ala	ND	40	ND	18
<i>gltBD argP-S94L</i>	Lys	ND	107	ND	44
<i>gltBD argP-S94L</i>	Lys-Ala	ND	111	ND	52

^a Intracellular levels of glutamate and glutamine were measured in cells that were grown as described in the text and were harvested from (i) glucose-minimal A medium with glycine betaine (low osmolarity) or (ii) glucose-minimal A medium with glycine betaine and 0.6 M NaCl (high osmolarity) without any additional supplement (None) or with Lys or Lys-Ala. Each value is the average of at least two independent determinations, and the variation between individual measurements was less than 20%.

^b The strains employed were wild-type strain MC4100, *gltBD* strain GJ4652, *proU gltBD* strain GJ2560, *argP gltBD* strain GJ4654, *glnE gltBD* strain GJ4663, and strain GJ4652/pHYD953 with the dominant *argP* mutation (*gltBD argP-S94L*).

^c ND, not determined.




FIG. 3. Dominant *argP-S94L* (*argP^Δ*) mutation and Lys effects on osmosensitivity in derivatives of *gltBD* strain GJ4652. Semilogarithmic growth curves were plotted for pBR329 (vector) (● and ○) and pHYD953 (*argP^Δ*) (▼ and ▽) derivatives of GJ4652 after 1:100 inoculation (from cultures grown in the stationary phase in glucose-minimal A medium) into glucose-minimal A medium containing 0.75 M NaCl, glycine betaine, and ampicillin without (● and ▼) or with (○ and ▽) Lys.

gltBD derivative was markedly increased (nearly 10-fold) compared with that in MC4100. Our results therefore indicate that osmosensitivity in the *argP gltBD* and *glnE gltBD* mutants (but not in the *proU gltBD* strain) is correlated with a failure to accumulate sufficient glutamate at an elevated osmolarity.

Exogenous Lys (or Lys-Ala) inhibits NH₄⁺ assimilation and confers osmosensitivity in *gltBD* mutant. A recent study in our laboratory (34) showed (i) that ArgP is a transcriptional activator of the *argO* gene (previously called *yggA*) that encodes a putative arginine exporter in *E. coli*; (ii) that intracellular Lys accumulation apparently correlates with abolition of the activator function of ArgP, as evidenced by the shutting down of *argO* transcription in *argP⁺* strains grown in medium supplemented with either Lys or the dipeptide Lys-Ala (which releases Lys by hydrolysis within the cells after uptake from the medium and hence serves to distinguish the [indirect] extracellular effects of Lys from the [direct] intracellular effects); and (iii) that a dominant gain-of-function *argP* mutation (*argP-S94L*), predicted to cause a Ser-to-Leu substitution at residue 94 of the encoded protein, renders *argO* expression high and constitutive with respect to Lys or Lys-Ala supplementation.

In light of the findings in the present study that an *argP* null mutation confers both osmosensitivity and an NH₄⁺ assimilation deficiency in the *gltBD* background, the effects of exogenous Lys or Lys-Ala supplementation on the growth of the *argP⁺ gltBD* strain were examined. We found that Lys addition inhibited growth of the *gltBD* mutant, both in medium with 2 mM NH₄⁺ as a nitrogen source (as shown in Fig. 2 for a *gltBD* derivative carrying the plasmid vector pBR329) and in a high-osmolarity medium (Fig. 3), just as an *argP* null mutation did.

Inhibition by Lys of both NH_4^+ assimilation and osmotolerance was also observed in a *gltBD* derivative carrying *argP*⁺ on a multicopy plasmid (data not shown). Lys supplementation did not further exacerbate the same phenotypes in an *argP* *gltBD* double mutant (data not shown).

The effects of exogenous Lys on the *gltBD* mutant were also mimicked by the effects of the dipeptide Lys-Ala for both inhibition of NH_4^+ assimilation (Fig. 2) and osmosensitivity (data not shown), whereas a control dipeptide histidylalanine had no effect (data not shown). Supplementation with Lys or Lys-Ala was also associated with a reduction in intracellular glutamate accumulation in high-osmolarity-grown cultures of the *gltBD* single mutant, to much the same extent that was observed upon introduction of the *argP* null mutation (Table 5).

Suppression by the dominant gain-of-function *argP*-S94L mutation of Lys and Lys-Ala effects in a *gltBD* strain. The inhibitory effects of Lys and Lys-Ala on *argO* transcription are abolished in strains with a dominant constitutive *argP*-S94L mutation (34). Accordingly, we tested whether the inhibition by these two compounds of NH_4^+ assimilation and osmotolerance in *gltBD* strains was affected by the dominant *argP*-S94L mutation. Growth comparisons between a pair of isogenic *gltBD* derivatives, one carrying plasmid pHYD953 with the dominant *argP*-S94L mutation (test strain) and the other carrying plasmid vector pBR329 (control strain), demonstrated that the dominant *argP*-S94L mutant allele conferred insensitivity to Lys or Lys-Ala supplementation on low- NH_4^+ medium (Fig. 2). The test strain was also insensitive to the inhibitory effect of Lys in high-osmolarity growth medium and indeed was somewhat more osmotolerant even in presence of Lys than the control strain was in its absence (Fig. 3); likewise, osmosensitivity in the *gltBD* strain associated with Lys-Ala supplementation was suppressed by the dominant *argP*-S94L mutation (data not shown). Concomitantly, the accumulation of cytoplasmic glutamate at high osmolarity was not affected by Lys or Lys-Ala supplementation in the *gltBD* *argP* derivative with the dominant *argP* mutation, unlike the accumulation of cytoplasmic glutamate in the *gltBD* *argP*⁺ strain (Table 5). These results suggest that the inhibitory effect of intracellular Lys on growth of the *gltBD* strain in either low- NH_4^+ or high-osmolarity medium, like the effect on *argO* expression, is mediated by the product of the *argP*⁺ gene.

***argO* is not involved in osmoregulation.** Given the parallels between the effects of *argP* (and the dominant *argP* mutation) on transcriptional regulation of the *argO*-encoded arginine exporter on the one hand (34) and on NH_4^+ assimilation and osmotolerance of *gltBD* strains on the other hand, we performed experiments to test whether the latter effect is in some way mediated by the former. Null mutants with mutations in *argO*, unlike mutants with mutations in *argP*, exhibited neither osmosensitivity nor NH_4^+ assimilation deficiency in the *gltBD* background (data not shown). Suppression by the dominant *argP*-S94L mutation of the inhibitory effect of Lys supplementation on osmotolerance and NH_4^+ assimilation in *gltBD* strains was not affected by introduction of an *argO* insertion mutation (data not shown). These results are consistent with the hypothesis (see below) that the *gltBD*-related phenotypes reflect ArgP-mediated regulation of a gene different from *argO*

that is involved in NH_4^+ assimilation through a GOGAT-independent pathway.

Additional evidence for the putative dual role of *argP*, for regulation of *argO* on the one hand and for regulation of a gene(s) involved in osmoregulation on the other, was obtained from an activator titration experiment. When the multicopy test plasmid pHYD954 with the cloned *cis* regulatory region of *argO* or the control plasmid vector pBluescriptII-KS was introduced into the *argP*⁺ *gltBD* strain GJ4652, derivatives with the former plasmid but not derivatives with the latter plasmid exhibited both osmosensitivity and a deficiency in NH_4^+ assimilation comparable to the results obtained with the *argP* *gltBD* derivatives described above (data not shown). Our interpretation is that the *argO* regulatory region in multiple copies titrates the ArgP protein so that this protein is not available for activation of the gene involved in NH_4^+ assimilation.

DISCUSSION

GOGAT-deficient (*gltBD*) mutants have previously been shown to be osmosensitive in medium with limiting NH_4^+ (14, 43). In this study, starting from a population of transposon-mutagenized cells of a *gltBD* strain, we identified *E. coli* derivatives with null mutations in *proU*, *argP*, or *glnE* that were osmosensitive in NH_4^+ -replete medium. In all these mutants, the absence of GOGAT was necessary for osmosensitivity.

The *proU* and *proP* loci encode active uptake systems that mediate the osmoprotectant effects of glycine betaine and proline in the enterobacteria (12, 13). The two transporters have overlapping functions, so that loss of either one is not associated with significant osmosensitivity (11, 17, 18, 24). Therefore, the isolation in the present study of osmosensitive *proU* insertion mutants in the *gltBD* background is a validation of the hypothesis that a GOGAT-deficient strain is sensitive to mutations affecting osmoregulation; Cayley et al. (5) have shown that glycine betaine accumulation in high-osmolarity-grown cells of a *proU*⁺ *proP*⁺ strain is associated with a reduction in intracellular glutamate levels.

Insertions at the other two loci, *glnE* and *argP*, also conferred osmosensitivity only in the GOGAT-deficient strain. We propose below that each of these genes is synergistic (for different reasons) with *gltBD* in reducing intracellular glutamate accumulation under osmotic stress conditions. Additional experimental support for the notion of synergism with the *gltBD* mutation has come from our findings (data not shown) that (i) in selection in transduction for osmotolerant derivatives of the *glnE* *gltBD* and *argP* *gltBD* strains, a large proportion of the transductants in both crosses had become *gltBD*⁺; and (ii) osmotolerance in both the *argP* *gltBD* and *glnE* *gltBD* mutants was fully restored in medium supplemented with aspartate, which serves a bypass route for glutamate synthesis. Therefore, it may be concluded that the strains are osmosensitive because they are limited for glutamate as a compatible solute and counterion for K^+ . (The alternative possibility, that glutamate is limiting for protein synthesis in these strains, is unlikely given the data shown in Table 5 for substantial intracellular pools of the amino acid, unless one assumes that the K_m for charging of glutamyl-tRNA is increased at an elevated osmolarity.) It may also be noted that although glycine betaine is known to mediate a K^+ - and glutamate-sparing effect in osmoregulation (5,

26), the NaCl concentrations used in this study were high enough to override this effect.

Osmosensitivity of *glnE* *gltBD* mutants. Whereas in a wild-type strain, increased activity of GS permits (and indeed is required for) growth on low levels of NH_4^+ or alternative nitrogen sources (35–37), excessive GS activity (particularly in a GOGAT-deficient strain) traps the assimilated nitrogen in glutamine. Adenylation by the *glnE*-encoded adenylyltransferase is one mechanism by which the activity of GS is down-regulated, but *glnE* mutants are nevertheless phenotypically almost normal because of the presence of additional mechanisms for *glnA* transcriptional regulation. Accordingly, *glnE* mutants have previously been shown to exhibit a growth deficiency only (i) when GS is constitutively expressed because of a *glnA* promoter mutation (25) or (ii) transiently upon shift-up from nitrogen-poor to high- NH_4^+ conditions (25, 43).

The synergism that was observed in this study between *glnE* and *gltBD* in conferring osmosensitivity may be explained by the following model: (i) during steady-state growth of the wild-type strain in NH_4^+ -replete media, the adenylation of GS (which is synthesized only at a basal level) is dispensable at low osmolarity but is required at high osmolarity in order to prevent the channeling of the accumulated glutamate into glutamine; and (ii) in the *glnE* *gltBD*⁺ strain subjected to osmotic stress, GOGAT is able to catalyze the conversion of glutamine to glutamate. Goss et al. (16) similarly explained the inability of *gltBD* mutants to utilize as nitrogen sources even compounds, such as proline, that can be catabolized to yield glutamate (37), on the grounds that GOGAT is needed to enable reconversion of glutamine to glutamate. Our model is consistent with the data in Table 5 which show that, compared with the wild-type strain or even the *gltBD* single mutant, the *glnE* *gltBD* double mutant had a reduced glutamate pool and a greatly increased glutamine pool after growth in the high-osmolarity medium.

Another implication of our model is that intracellular glutamine cannot substitute for glutamate in osmoregulation. This conclusion is consistent with the findings obtained in a previous study (14), as well as with the hypothesis that glutamate's primary role in osmoregulation is to serve as the counterion to K^+ (12, 13).

Osmosensitivity of *argP* *gltBD* mutants. Null mutations in *argP*, unlike null mutations in *glnE*, were associated with a reduction in the ability of the *gltBD* mutant to grow in medium with a low NH_4^+ concentration. Compared with the *gltBD* single mutant, the *argP* *gltBD* strain exhibited reduced GDH activity in NH_4^+ -replete medium, particularly at elevated osmolarity (Table 4). Thus, we concluded that the GDH-catalyzed pathway of NH_4^+ assimilation is compromised in the *argP* mutants and that the reduced capacity for glutamate synthesis in the *argP* *gltBD* strains may be sufficient for growth in low-osmolarity media but not for growth in high-osmolarity media. A defect in glutamate accumulation in the double mutant was directly demonstrated in this study (Table 5). Whether an NH_4^+ uptake defect also might contribute to the *argP* mutant phenotypes remains to be determined.

A previous study (39) showed that *fnr* mutations also reduce NH_4^+ assimilation and confer osmosensitivity in the *gltBD* background and that multiple copies of *spoT*⁺ reverse both phenotypes in an *fnr* *gltBD* strain. We found in this work that

multiple copies of *spoT*⁺ did not affect either phenotype in an *argP* *gltBD* strain (data not shown).

The mechanism by which a null mutation in *argP* reduces GDH activity, and consequently NH_4^+ assimilation and osmotolerance, in the *gltBD* strain is not known. Evidence from a study recently completed in our laboratory (34) suggests that ArgP is ordinarily a transcriptional activator protein for a target gene, *argO* (involved in arginine efflux), and that in the presence of intracellular Lys as a coeffector it represses *argO* expression. Thus, the results described here for the effects of *argP* mutations and of Lys or Lys-Ala supplementation in *gltBD* strains are most simply explained by the hypothesis that there is another gene (which perhaps is *gdhA* itself) whose regulation mirrors that of *argO*, which is involved in determining or modulating GDH activity in the *gltBD* strain. The hypothesis that intracellular Lys likely phenocopies the *argP* null mutation for decreased glutamate accumulation and osmosensitivity is strongly supported by our observation that the effects of exogenous Lys or Lys-Ala are suppressed in strains carrying a Lys-insensitive dominant gain-of-function *argP* allele.

Interestingly, Bender and coworkers have shown that *gdhA* transcription in *Klebsiella aerogenes* is repressed about threefold upon Lys supplementation; to explain this, these authors postulated that an unidentified regulator protein which is Lys sensitive activates *gdhA* in the strain (15, 22). Based on our results, it is likely that this regulator protein is the ortholog of *E. coli* ArgP.

ACKNOWLEDGMENTS

We thank S. Blanquet, E. Bremer, K. Isono, and C. G. Lerner for providing strains, phages, and plasmids. We acknowledge V. Vamsee Krishna and T. Giri Babu for technical assistance, Mehar Sultana for primer synthesis, N. Nagesh for DNA sequencing, S. Jamal Mohamed Nurul Jain and S. M. Naushad for help with reverse-phase high-performance liquid chromatography analysis, and members of the laboratory of J.G. for advice and discussions.

This work was supported by research fellowships to M.R.N. and R.S.L. from the Council of Scientific and Industrial Research and by a grant-in-aid from the Department of Biotechnology, Government of India (project BT/PR2430). J.G. is an honorary faculty member of the Jawaharlal Nehru Centre for Advanced Scientific Research.

REFERENCES

- Berlyn, M. K. B. 1998. Linkage map of *Escherichia coli* K-12, edition 10: the traditional map. *Microbiol. Mol. Biol. Rev.* **62**:814–984.
- Blattner, F. R., G. Plunkett III, C. A. Bloch, N. T. Perna, V. Burland, M. Riley, J. Collado-Vides, J. D. Glasner, C. K. Rode, G. F. Mayhew, J. Gregor, N. W. Davis, H. A. Kirkpatrick, M. A. Goeden, D. J. Rose, B. Mau, and Y. Shao. 1997. The complete genome sequence of *Escherichia coli* K-12. *Science* **277**:1453–1462.
- Botsford, J. L., M. Alvarez, R. Hernandez, and R. Nicholas. 1994. Accumulation of glutamate by *Salmonella typhimurium* in response to osmotic stress. *Appl. Environ. Microbiol.* **60**:2568–2574.
- Bradford, M. M. 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* **72**:248–254.
- Cayley, S., B. A. Lewis, and M. T. Record, Jr. 1992. Origins of the osmoprotective properties of betaine and proline in *Escherichia coli* K-12. *J. Bacteriol.* **174**:1586–1595.
- Celis, R. T. F. 1990. Mutant of *Escherichia coli* K-12 with defective phosphorylation of two periplasmic transport proteins. *J. Biol. Chem.* **265**:1787–1793.
- Celis, R. T. F. 1999. Repression and activation of arginine transport genes in *Escherichia coli* K-12 by the ArgP protein. *J. Mol. Biol.* **294**:1087–1095.
- Celis, R. T. F., H. J. Rosenfeld, and W. K. Maas. 1973. Mutant of *Escherichia coli* K-12 defective in the transport of basic amino acids. *J. Bacteriol.* **116**:619–626.
- Chang, A. C., and S. N. Cohen. 1978. Construction and characterization of amplifiable multicopy DNA cloning vehicles derived from the P15A cryptic mini-plasmid. *J. Bacteriol.* **134**:1141–1156.

10. Covarrubias, L., and F. Bolivar. 1982. Construction and characterization of new cloning vehicles. VI. Plasmid pBR329, a new derivative of pBR328 lacking the 482-base-pair inverted duplication. *Gene* **17**:79–89.
11. Csonka, L. N. 1982. A third L-proline permease in *Salmonella typhimurium* which functions in media of elevated osmotic strength. *J. Bacteriol.* **151**:1433–1443.
12. Csonka, L. N. 1989. Physiological and genetic responses of bacteria to osmotic stress. *Microbiol. Rev.* **53**:121–147.
13. Csonka, L. N., and W. Epstein. 1996. Osmoregulation, p. 1210–1223. In F. C. Neidhardt, R. Curtiss III, J. L. Ingraham, E. C. C. Lin, K. B. Low, B. Magasanik, W. S. Reznikoff, M. Riley, M. Schaechter, and H. E. Umbarger (ed.), *Escherichia coli* and *Salmonella*: cellular and molecular biology, 2nd ed. American Society for Microbiology, Washington, D.C.
14. Csonka, L. N., T. P. Ikeda, S. A. Fletcher, and S. Kustu. 1994. The accumulation of glutamate is necessary for optimal growth of *Salmonella typhimurium* in media of high osmolality but not induction of the *proU* operon. *J. Bacteriol.* **176**:6324–6333.
15. Goss, T. J., B. K. Janes, and R. A. Bender. 2002. Repression of glutamate dehydrogenase formation in *Klebsiella aerogenes* requires two binding sites for the nitrogen assimilation control protein, NAC. *J. Bacteriol.* **184**:6966–6975.
16. Goss, T. J., A. Perez-Matos, and R. A. Bender. 2001. Roles of glutamate synthase, *gltBD*, and *gltF* in nitrogen metabolism of *Escherichia coli* and *Klebsiella aerogenes*. *J. Bacteriol.* **183**:6607–6619.
17. Gowrishankar, J. 1985. Identification of osmoreponsive genes in *Escherichia coli*: evidence for participation of potassium and proline transport systems in osmoregulation. *J. Bacteriol.* **164**:434–445.
18. Gowrishankar, J. 1986. *proP*-mediated proline transport also plays a role in *Escherichia coli* osmoregulation. *J. Bacteriol.* **166**:331–333.
19. Gowrishankar, J., and J. Pittard. 1982. Construction from Mu d1 (*lac Ap*^r) lysogens of lambda bacteriophage bearing promoter-*lac* fusions: isolation of λ *pphA-lac*. *J. Bacteriol.* **150**:1122–1129.
20. Hwang, D. S., and A. Kornberg. 1990. A novel protein binds a key origin sequence to block replication of an *E. coli* minichromosome. *Cell* **63**:325–331.
21. Hwang, D. S., B. Thony, and A. Kornberg. 1992. IciA protein, a specific inhibitor of initiation of *Escherichia coli* chromosomal replication. *J. Biol. Chem.* **267**:2209–2213.
22. Janes, B. K., P. J. Pomposiello, A. Perez-Matos, D. J. Najarian, T. J. Goss, and R. A. Bender. 2001. Growth inhibition caused by overexpression of the structural gene for glutamate dehydrogenase (*gdhA*) from *Klebsiella aerogenes*. *J. Bacteriol.* **183**:2709–2714.
23. Kohara, Y., K. Akiyama, and K. Isono. 1987. The physical map of the whole *E. coli* chromosome: application of a new strategy for rapid analysis and sorting of a large genomic library. *Cell* **50**:495–508.
24. Koo, S. P., and I. R. Booth. 1994. Quantitative analysis of growth stimulation by glycine betaine in *Salmonella typhimurium*. *Microbiology* **140**:617–621.
25. Kustu, S., J. Hirschman, D. Burton, J. Jelesko, and J. C. Meeks. 1984. Covalent modification of bacterial glutamine synthetase: physiological significance. *Mol. Gen. Genet.* **197**:309–317.
26. Larsen, P. I., L. K. Sydnes, B. Landfald, and A. R. Strom. 1987. Osmoregulation in *Escherichia coli* by accumulation of organic osmolytes: betaines, glutamic acid, and trehalose. *Arch. Microbiol.* **147**:1–7.
27. Lerner, C. G., and M. Inouye. 1990. Low copy number plasmids for regulated low-level expression of cloned genes in *Escherichia coli* with blue/white insert screening capability. *Nucleic Acids Res.* **18**:4631.
28. Leveque, F., S. Blanchin-Roland, G. Fayat, P. Plateau, and S. Blanquet. 1990. Design and characterization of *Escherichia coli* mutants devoid of Ap₄N-hydrolase activity. *J. Mol. Biol.* **212**:319–329.
29. May, G., E. Faatz, M. Villarejo, and E. Bremer. 1986. Binding protein dependent transport of glycine betaine and its osmotic regulation in *Escherichia coli* K12. *Mol. Gen. Genet.* **205**:225–233.
30. McLaggan, D., T. M. Logan, D. G. Lynn, and W. Epstein. 1990. Involvement of γ -glutamyl peptides in osmoadaptation of *Escherichia coli*. *J. Bacteriol.* **172**:3631–3636.
31. McLaggan, D., J. Naprstek, E. T. Buurman, and W. Epstein. 1994. Interdependence of K⁺ and glutamate accumulation during osmotic adaptation of *Escherichia coli*. *J. Biol. Chem.* **269**:1911–1917.
32. Meers, J. L., D. W. Tempest, and C. M. Brown. 1970. 'Glutamine (amide): 2-oxoglutarate amino transferase oxidoreductase (NADP)', an enzyme involved in the synthesis of glutamate by some bacteria. *J. Gen. Microbiol.* **64**:187–194.
33. Miller, J. H. 1992. A short course in bacterial genetics: a laboratory manual and handbook for *Escherichia coli* and related bacteria. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
34. Nandineni, M. R., and J. Gowrishankar. 2004. Evidence for an arginine exporter encoded by *yggA* (*argO*) that is regulated by the LysR-type transcriptional regulator ArgP in *Escherichia coli*. *J. Bacteriol.* **186**:3539–3546.
35. Reitzer, L. 2003. Nitrogen assimilation and global regulation in *Escherichia coli*. *Annu. Rev. Microbiol.* **57**:155–176.
36. Reitzer, L., and B. L. Schneider. 2001. Metabolic context and possible physiological themes of σ^{54} -dependent genes in *Escherichia coli*. *Microbiol. Mol. Biol. Rev.* **65**:422–444.
37. Reitzer, L. J. 1996. Ammonia assimilation and the biosynthesis of glutamine, glutamate, aspartate, asparagine, L-alanine, and D-alanine, p. 391–407. In F. C. Neidhardt, R. Curtiss III, J. L. Ingraham, E. C. C. Lin, K. B. Low, B. Magasanik, W. S. Reznikoff, M. Riley, M. Schaechter, and H. E. Umbarger (ed.), *Escherichia coli* and *Salmonella*: cellular and molecular biology, 2nd ed. American Society for Microbiology, Washington, D.C.
38. Sambrook, J., E. F. Fritsch, and T. Maniatis. 1989. Molecular cloning: a laboratory manual, 2nd ed. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
39. Saroja, G. N., and J. Gowrishankar. 1996. Roles of SpoT and FNR in NH₄⁺ assimilation and osmoregulation in GOGAT (glutamate synthase)-deficient mutants of *Escherichia coli*. *J. Bacteriol.* **178**:4105–4114.
40. Smith, G. R., Y. S. Halpern, and B. Magasanik. 1971. Genetic and metabolic control of enzymes responsible for histidine degradation in *Salmonella typhimurium*. *J. Biol. Chem.* **246**:3320–3329.
41. Spiro, S., and J. R. Guest. 1988. Inactivation of the FNR protein of *Escherichia coli* by targeted mutagenesis in the N-terminal region. *Mol. Microbiol.* **2**:701–707.
42. Thony, B., D. S. Hwang, L. Fradkin, and A. Kornberg. 1991. *iciA*, an *Escherichia coli* gene encoding a specific inhibitor of chromosomal initiation of replication *in vitro*. *Proc. Natl. Acad. Sci. USA* **88**:4066–4070.
43. Yan, D., T. P. Ikeda, A. E. Shauger, and S. Kustu. 1996. Glutamate is required to maintain the steady-state potassium pool in *Salmonella typhimurium*. *Proc. Natl. Acad. Sci. USA* **93**:6527–6531.