

PERMANENT GENETIC RESOURCES NOTE

Polymorphic microsatellite loci for primitively eusocial wasp *Ropalidia marginata*

S. JOHNY,*† S. CHAKRABORTY,*† R. GADAGKAR* and J. NAGARAJU†

*Centre for Ecological Sciences, Indian Institute of Science, Bangalore 560012, India, †Laboratory of Molecular Genetics, Centre for DNA Fingerprinting and Diagnostics, ECIL Road, Nacharam, Hyderabad 500 076, India

Abstract

We report here development and characterization of 48 novel microsatellite markers for *Ropalidia marginata*, a tropical, primitively eusocial polistine wasp from peninsular India. Thirty-two microsatellites showed polymorphism in a wild population of *R. marginata* (N = 38) collected from Bangalore, India. These markers will facilitate answering some interesting questions in ecology and evolutionary biology of this wasp, such as population structure, serial polygyny, intra-colony genetic relatedness and the pattern of queen succession.

Keywords: genetic variation, microsatellites, queen succession, *Ropalidia marginata*

Received 5 June 2008; accepted 21 January 2009

Ropalidia marginata (Lep.) (Hymenoptera: Vespidae) is a tropical polistine wasp, widely distributed in peninsular India (Gadagkar 2001). At a certain point of time, a nest consists of only one queen and several workers, although a nest can be under the regime of several queens during its existence – a phenomenon known as serial polygyny. This species is considered primitively eusocial, because of the absence of morphological differentiation between queens and workers and because many, if not all, female wasps can mate, develop their ovaries and function as solitary nest foundresses or as queens of multiple foundress nests (Gadagkar 2001). However, there is growing evidence that *R. marginata* is different from other primitively eusocial species. In all other primitively eusocial species studied so far, queens are the behaviourally most dominant and active individuals on the nest, and thus use physical aggression (dominance behaviour) to suppress worker reproduction. In contrast, *R. marginata* queens are described as behaviourally nondominant, docile sitters, who cannot possibly inhibit worker reproduction by physical aggression and who probably use pheromones to do so (Sumana & Gadagkar 2003). Because of the wealth of behavioural information already available on *R. marginata* (Kardile & Gadagkar 2003; Sumana & Gadagkar 2003; Agrahari & Gadagkar 2004; Bruyndonckx *et al.* 2006; Bhadra *et al.* 2007), the availability of molecular markers will further enhance our understanding of this species, especially the role of

intra-colony genetic relatedness in social evolution and the pattern of queen succession. Therefore, in the present study, we have developed informative microsatellite markers for *R. marginata*. These markers can also be employed in studying population genetic structure of this species.

In order to develop microsatellite markers, repeat enriched genomic library was constructed using the following procedure. Genomic DNA was extracted from adults of *R. marginata* collected from the Indian Institute of Science campus, Bangalore, India. A pool of 20 adults from different nests was collected and used for DNA extraction. Hybridization capture approach (Glenn & Schable 2005) was employed for enriching the DNA pool with repeat harbouring fragments. In brief, genomic DNA was digested with the restriction enzyme *Rsa*I (New England Biolabs, UK), and 300–1000-bp inserts were ligated into the super SNX double-stranded linker on both sides. The DNA was hybridized with biotinylated oligonucleotides such as (ATT)₈, (ATGA)₇, (CA)₁₀, (GA)₁₀, (GATA)₇ and (CAC)₇, and separated with Dynabeads (Dyna) under magnetic field. The products eluted from beads were polymerase chain reaction (PCR) amplified using the super SNX forward primer (GTTTAAGGCCTAGCTAGCAGAATC) and cloned into pCR2.1-TOPO vector (Invitrogen). Resulted recombinant plasmids were transformed into XL1-Blue competent cells. Plasmids were extracted from clones with more than 200-bp inserts, using QIAprep Spin Miniprep Kit (QIAGEN). The inserts were sequenced using BigDye terminator

Correspondence: J. Nagaraju, E-mail: jnagaraju@cdfd.org.in

2 PERMANENT GENETIC RESOURCES NOTE

Table 1 Characteristics of *Ropalidia marginata* microsatellite loci genotyped on ABI PRISM 3730 DNA analyser, including repeat motif, primer sequences, annealing temperature, number of individuals screened (*N*), number of alleles (*N_a*), observed (*H_O*) and expected heterozygosities (*H_E*). Data obtained upon screening 38 individuals sampled from three locations of Bangalore region

Locus (GenBank)	Fluorescent label	Colour	Repeat motif	Primer	Ann. temp.	<i>N/N_a</i>	Size range	<i>H_O</i>	<i>H_E</i>
RmSat006† (FJ589173)	TAM	Yellow	(TAAAAA) ₃	F: TACCGCAACATTTTATTTC R: TCCTGTCTCAATACCGTTAGA	56 °C	36/5	266–274	0.194	0.603*
RmSat010† (FJ589175)	JOE	Green	(CCA) ₂ A(CCA) ₇ (GCA) ₂ ... (CT) ₅	F: GAATCTTCTCTCATCCACCA R: GTGTAGCATCCCTCGTAAAG	56 °C	32/10	132–145	0.531	0.791*
RmSat011† (FJ589176)	TAM	Yellow	(TAAA) ₉	F: ATGCGTTTCCGACCTGAGCT R: CGAATTCCTTAGGCTTATAG	56 °C	36/8	226–247	0.361	0.749*
RmSat013 (FJ589178)	JOE	Green	(TAA) ₃ TCA(TAA) ₃ TCA(TAA) ₃	F: TAGGAACCGAATCTACCAAC R: CAAATCCGATTTGTGACGTAA	58 °C	34/10	106–117	0.706	0.791*
RmSat019 (FJ589179)	FAM	Blue	(CA) ₅ (CAAA) ₂ (CA) ₂ ... (CATA) ₂ (CA) ₁₁	F: TCTCAGTTTCATTTCGATTT R: CTTTTTCGTGATATCGACTGC	56 °C	32/14	233–256	0.781	0.895*
RmSat028† (FJ589182)	FAM	Blue	(CT) ₉ (CTTT) ₃	F: GTGCAGCCCAACCGATACT R: TCCTCCCGGATATCACATA	58 °C	35/11	247–264	0.571	0.822*
RmSat065† (FJ589193)	TAM	Yellow	(CTGA) ₂ (CTTA) ₄ ... (CA) ₁₃ (CAGC) ₄	F: TTCAAGGACAAACGCTATTT R: GTTTGAACGTAAGGGAGTG	56 °C	35/16	162–186	0.543	0.928*
RmSat067† (FJ589194)	JOE	Green	(CAA) ₆	F: CACTTCTCTCCACTTGAG R: CGTGGATAAAGATTGGTGAT	55 °C	35/6	221–228	0.0571	0.661*
RmSat074 (FJ589197)	TAM	Yellow	(GTGA) ₈ ... (GA) ₅	F: AGAGAAGGAAAGGGACAGAG R: CTAACAGCAACAGCACTCCT	56 °C	30/8	101–133	0.700	0.820
RmSat083† (FJ589201)	JOE	Green	(AGT) ₇ ... (AGT) ₁₀	F: GCCAACAGATCCAACAGTAA R: CCTGTTCTACTGCTGCTAC	56 °C	37/16	170–191	0.514	0.893*

*Significant deviation from Hardy–Weinberg equilibrium. Loci marked (†) denote presence of null alleles.

chemistry on an ABI 3100 automated sequencer (Applied Biosystems) with M13 primers.

Of the 415 clones of *R. marginata* microsatellite-enriched library, 148 were picked and their plasmid inserts were sequenced. The sequences were screened for the presence of microsatellites using the software Micas (www.cdfd.org.in/micas). Primers were designed for sequences flanking the microsatellites using Primer 3 software (Rozen & Skaletsky 2000). Each primer set was tested for proper amplification by PCR. Optimal annealing temperature for each set was obtained using a temperature gradient thermal cycler (Eppendorf GmbH).

Primers were designed for 84 repeat-containing loci and 48 were selected for subsequent screening for polymorphism. The others were rejected due to nonspecific amplification or absence of PCR product. Among these microsatellite loci, the majority of them were dinucleotide repeat motifs (66%) followed by tri- (26%) and tetranucleotide repeat motifs. A very low percentage of hexa- and heptanucleotide repeat motifs (2% each) were also identified in this study. Among the dinucleotide repeats, the GA repeat motif was the most common accounting for 69% followed by 27% of GT repeat motifs, suggesting that the GA/GT repeats are abundant in hymenoptera as reported by others (Krieger & Keller 1997; Gopinath *et al.* 2001; Paxton *et al.* 2003; Solignac *et al.* 2003; Arthofer *et al.* 2005). The length of dinucleotide repeats varied from five to 28, as compared to

trinucleotide repeats which varied between four and 13. For the population genetic study, wasps sampled were preserved in 70% ethyl alcohol and DNA was extracted using HotSHOT (Truett *et al.* 2000) method at the earliest opportunity.

Fourteen microsatellite loci were confirmed to be monomorphic and 32 loci were found to be polymorphic. Two loci namely RmSat043 and RmSat070 were highly polymorphic but produced unscorable stutter bands.

Out of 32 polymorphic loci, 10 were evaluated using fluorescent-labelled primers on an ABI PRISM 3730 DNA analyser using DNA from 38 females sampled from three different localities across Bangalore, India (Table 1). The remaining 22 loci were evaluated on 12% PAGE using DNA from 38 female wasps sampled from another population collected from Bangalore (Table 2). The primers were labelled using three different fluorescent dyes, that is, TAM (Yellow), JOE (Green) and FAM (Blue). The PCR was performed on an Applied Biosystems, GeneAmp PCR System 9700. The amplification conditions include an initial denaturation at 94 °C for 3 min, 30 cycles of 94 °C for 30 s, a locus-specific annealing temperature (Tables 1 and 2) and 72 °C for 45 s, followed by a final extension of 72 °C for 10 min. The 10 loci given in Table 1 were run on an ABI PRISM 3730 DNA analyser, with Pop-7 as sieving matrix, HiDi Formamide as single-stranded DNA stabilizer and GeneScan 500 ROX standard as a size marker. ABI

Table 2 Characteristics of *Ropalidia marginata* microsatellite loci on polyacrylamide gel, including repeat motif, primer sequences, annealing temperature, number of individuals screened (N), number of alleles (N_a), observed (H_o) and expected heterozygosities (H_e). Data obtained upon screening 38 individuals sampled from a population collected from Bangalore region

Locus	Repeat motif	Primer	Ann. temp.	N/N_a	Size range	H_o	H_e
RmSat001 (FJ589170)	GCT(ACT) ₂ (GCT) ₁₁	F: ATTTCGAACAAAAGCACCT R: AGGGATAACCGAAATGGA	58 °C	38/3	273–300	0.395	0.442
RmSat002† (FJ589171)	(CA) ₉ ... (CAG) ₆ (CTG) ₂	F: GCGACTCAATTTCTGTTTCT R: AAACCAATAACAGCAACCAC	58 °C	38/6	187–210	0.378	0.783*
RmSat005† (FJ589172)	(TA) ₅	F: TAAAAAGGGATGGATGTGTC R: TCATCGGACAAACTGTGAG	58 °C	38/3	146–156	0.211	0.482*
RmSat007 (FJ589174)	(CA) ₅ ... (GCACCA) (GAACCA) ₃ (GCACCA)	F: CAAGTGACATGTCGATGAGA R: AGTAAACGGAGCTACTGCTG	58 °C	38/2	217, 229	0.316	0.438
RmSat012† (FJ589177)	(CGT) ₂ TGT(CGT) ₁₃	F: TGTTCATCAAGTGACGAAACG R: CGTTGTTACATCGTTCAAAG	58 °C	37/7	113–149	0.605	0.760*
RmSat020† (FJ589180)	(CA) ₄ AG(CA) ₅ TA(CA) ₄ GA(CA) ₃	F: CGATCTAACTCTGTGGCTCT R: ACGTTCAGATTGAGTCCAAAT	58 °C	38/4	122–136	0.447	0.699*
RmSat022† (FJ589181)	(CA) ₁₉	F: CGATTAACGTAATTCATTATGG R: GTTGTGCTCTCGAAAAACT	56 °C	38/5	180–204	0.500	0.737*
RmSat029 (FJ589183)	(GA) ₂ AA(GA) ₄ AA(GA) ₃ AA(GA) ₃ AG(GA) ₂	F: GTATTGCATCGTGTGTTTGT R: AGAGCCACAGAGTTAGATCG	58 °C	38/2	117, 125	0.026	0.026
RmSat030 (FJ589184)	(GA) ₂ AA(GA) ₂₀	F: TCCCGTAGTTCCTATCAATG R: TTCATACACGTTTTTCATCCA	54 °C	38/5	136–154	1.000	0.771*
RmSat036 (FJ589185)	(CT) ₆ TT(CT) ₂	F: ACCAATAATACACACGCAGA R: TCGTATTTTCTCCCTCGATA	56 °C	38/2	222, 230	0.342	0.287
RmSat039 (FJ589186)	(GA) ₁₀	F: ATATGGGTTCGATGAAAAG R: ACATGAAAGACACACACGAA	56 °C	38/5	114–128	0.447	0.559*
RmSat041 (FJ589187)	(CT) ₇ C(CCT) ₂ T(CT) ₅	F: ACAACGCAATCTCCGTATTA R: GTTCAAGACGGTGAAAAATG	56 °C	38/2	165, 177	0.158	0.147
RmSat045† (FJ589188)	(GA) ₂ AA(GA) ₉ AA(GA) ₃ A(GA) ₂	F: CATAACTTCTACGCCCTTGT R: TAACGTTTCGCGATAAAGAC	56 °C	38/7	143–170	0.526	0.735*
RmSat047 (FJ589189)	(GGA) ₂ (GA) ₃ AA(GA) ₉	F: GATCGTCCCATACTCAGAGA R: CAGACCCGGGAGAATTTCTAT	56 °C	37/10	136–180	0.730	0.861*
RmSat049 (FJ589190)	(CT) ₈ ... (CT) ₁₀	F: TCACATAGACGCCAACACAT R: AGTCTAGGATGAGGGGAAG	56 °C	38/2	165 171	0.026	0.026
RmSat056 (FJ589191)	(GA) ₁₄	F: CGAACCATTCCACCACATAC R: CCGCGACCAACCAATAC	58 °C	38/8	88–132	0.789	0.837
RmSat064† (FJ589192)	(GT) ₁₀ AT(GT) ₄	F: AAAGAAGTTCGACCCAAAG R: ATTACAATTGCGAGCAAAC	54 °C	38/6	265–312	0.500	0.755*
RmSat071† (FJ589195)	(CT) ₄ CA(CT) ₁₃ TT(CT) ₃	F: CCATCACCCGAAGAACTAAT R: AGAAGACGTAGATACCCCAAC	55 °C	38/10	139–185	0.632	0.838*
RmSat072 (FJ589196)	(CT) ₁₁	F: CGACTAACTTCATCGTTTCT R: ACGCAACGATGTTTAATTTT	56 °C	38/2	147, 153	0.079	0.077
RmSat077† (FJ589198)	(GA) ₇ A(GAG) ₃	F: CTCTAAACCCCGATAATGTTT R: AACTAAACGGATGAAACGAA	56 °C	38/4	116–131	0.184	0.689*
RmSat079 (FJ589199)	(CT) ₉	F: CTGGTGCTAAGTAAAAATCC R: ATACGTGCGGTAATAAAT	56 °C	38/4	172–184	0.658	0.529
RmSat082 (FJ589200)	(CAA) ₆ (CAG) ₃ (CAA) ₂	F: TGTCCACAGTATCCTCAAT R: AATGACAGACTTTTGGGACT	56 °C	38/2	196, 202	0.053	0.052

*Significant deviation from Hardy–Weinberg equilibrium. Loci marked (†) denote presence of null alleles. Allele sizes for these loci may not be accurate as these loci were resolved on polyacrylamide gels.

PRISM GeneMapper software version 3.0 was used to size the alleles. The amplified products of the remaining 22 loci (Table 2) were separated on 12% polyacrylamide gels using Hoefer SE 600 standard vertical electrophoresis unit along with 50-bp DNA ladder (MBI Fermentas) and allelic

products were visualized by silver staining. The alleles were scored using Bio-Rad Quantity One software (Bio-Rad Laboratories). Analyses of polymorphism, including number of alleles, observed heterozygosity, expected heterozygosity, likelihood ratio test for linkage disequilibrium and exact

tests for the Hardy–Weinberg equilibrium were performed by using Arlequin version 3.1 (Excoffier *et al.* 2005). The probability of occurrence of null alleles was tested using Micro-Checker (van Oosterhout *et al.* 2004).

The number of alleles per locus ranged from two to 16, and the observed and expected heterozygosities ranged from 0.026 to 0.895. After sequential Bonferroni correction, 35 of the 45 combinations showed significant linkage disequilibrium in individuals scored on automatic DNA analyser. However, in case of the loci analysed on PAGE, none of the 231 pairwise comparisons between loci showed significant linkage disequilibrium. Significant deviations ($P < 0.05$) from Hardy–Weinberg equilibrium were detected in 21 out of 32 loci (Tables 1 and 2). Departure from Hardy–Weinberg equilibrium was probably due to heterozygote deficit. However, this could be interpreted as the result of Wahlund effect (Wahlund 1928), because the Bangalore population may have been structured into several subpopulations. Null alleles may be present in 16 of the 32 loci reported here. The informative microsatellite markers reported in the present study will be of great value for answering interesting questions in ecology and evolutionary biology of *R. marginata*.

Acknowledgements

Financial support to RG and JN by the Department of Biotechnology, New Delhi under the project 'Molecular Ecology and Systematics (Grant no. BT/012/01/NBDB/e2004)' is greatly acknowledged. SC was supported by a fellowship from Council for Scientific and Industrial Research, India during this work. K. P. Arunkumar helped with the data analysis.

References

Agrahari M, Gadagkar R (2004) Hard working nurses rather than over-aged nurses permit *Ropalidia marginata* to respond to the loss of young individuals. *Insectes Sociaux*, **51**, 306–307.

Arthofer W, Schlick-Steiner BC, Steiner FM *et al.* (2005) Isolation of polymorphic microsatellite loci for the study of habitat fragmentation in the harvester ant *Messor structor*. *Conservation Genetics* **6**, 859–861.

Bhadra A, Iyer PL, Sumana A *et al.* (2007) How do workers of the primitively eusocial wasp *Ropalidia marginata* detect the presence of their queens? *Journal of Theoretical Biology*, **246**, 574–582.

Bruyndonckx N, Kardile SP, Gadagkar R (2006) Dominance behaviour and regulation of foraging activity in the primitively eusocial wasp *Ropalidia marginata* (Lep.) (Hymenoptera: Vespidae). *Behavioural Processes*, **72**, 100–103.

Excoffier L, Laval G, Schneider S (2005) Arlequin ver. 3.0: an integrated software package for population genetics data analysis. *Evolutionary Bioinformatics Online*, **1**, 47–50.

Gadagkar R (2001) *The Social Biology of Ropalidia Marginata: Toward Understanding the Evolution of Eusociality*. Harvard University Press, Cambridge, Massachusetts.

Glenn TC, Schable NA (2005) Isolating microsatellite DNA loci. *Methods in Enzymology*, **395**, 202–222.

Gopinath A, Gadagkar R, Rao MRS (2001) Identification of polymorphic microsatellite loci in the queenless, ponerine ant *Diacamma ceylonense*. *Molecular Ecology Notes*, **1**, 126–127.

Kardile SP, Gadagkar R (2003) Regulation of worker activity in the primitively eusocial wasp *Ropalidia cyathiformis*. *Behaviour*, **140**, 1219–1234.

Krieger MJB, Keller L (1997) Polymorphism at dinucleotide microsatellite loci in fire ant *Solenopsis invicta* populations. *Molecular Ecology*, **6**, 997–999.

van Oosterhout CHW, Wills DPM, Shipley P (2004) Micro-Checker: software for identifying and correcting genotyping errors in microsatellite data. *Molecular Ecology Notes*, **4**, 536–538.

Paxton RJ, Arévalo E, Field J (2003) Microsatellite loci for the eusocial *Lasioglossum malachurum* and other sweat bees (Hymenoptera, Halictidae). *Molecular Ecology Notes*, **3**, 82–84.

Rozen S, Skaletsky H (2000) Primer 3 on the WWW for general users and for biologist programmers. *Methods in Molecular Biology*, **132**, 365–386.

Solignac M, Vautrin D, Loiseau A *et al.* (2003) Five hundred and fifty microsatellite markers for the study of the honeybee (*Apis mellifera* L.) genome. *Molecular Ecology Notes*, **3**, 307–311.

Sumana A, Gadagkar R (2003) *Ropalidia marginata* — a primitively eusocial wasp society headed by docile queens. *Current Science*, **84**, 1464–1468.

Truett GE, Heeger P, Mynatt RL, Truett AA, Walker JA, Warman ML (2000) Preparation of PCR-quality mouse genomic DNA with hot sodium hydroxide and tris (HotSHOT). *BioTechniques*, **29**, 52–54.

Wahlund S (1928) Zusammensetzung von Population und Korrelationserscheinung vom Standpunkt der Vererbungslehre aus betrachtet. *Hereditas*, **11**, 65–106.

Author Query Form

Journal: Molecular Ecology Resources

Article: mer_2597

During the copy-editing of your paper, the following queries arose. Please respond to these by marking up your proofs with the necessary changes/additions. Please write your answers on the query sheet if there is insufficient space on the page proofs. Please write clearly and follow the conventions shown on the attached corrections sheet. If returning the proof by fax do not write too close to the paper's edge. Please remember that illegible mark-ups may delay publication.

Many thanks for your assistance.

No.	Query	Remarks
1	Author: Please check if the URL for Micas is correct.	

MARKED PROOF

Please correct and return this set

Please use the proof correction marks shown below for all alterations and corrections. If you wish to return your proof by fax you should ensure that all amendments are written clearly in dark ink and are made well within the page margins.


<i>Instruction to printer</i>	<i>Textual mark</i>	<i>Marginal mark</i>
Leave unchanged	... under matter to remain	Ⓟ
Insert in text the matter indicated in the margin	∧	New matter followed by ∧ or ∧ [Ⓢ]
Delete	/ through single character, rule or underline or ┌───┐ through all characters to be deleted	Ⓞ or Ⓞ [Ⓢ]
Substitute character or substitute part of one or more word(s)	/ through letter or ┌───┐ through characters	new character / or new characters /
Change to italics	— under matter to be changed	↙
Change to capitals	≡ under matter to be changed	≡
Change to small capitals	≡ under matter to be changed	≡
Change to bold type	~ under matter to be changed	~
Change to bold italic	≈ under matter to be changed	≈
Change to lower case	Encircle matter to be changed	≡
Change italic to upright type	(As above)	⊕
Change bold to non-bold type	(As above)	⊖
Insert 'superior' character	/ through character or ∧ where required	Υ or Υ under character e.g. Υ or Υ
Insert 'inferior' character	(As above)	∧ over character e.g. ∧
Insert full stop	(As above)	⊙
Insert comma	(As above)	,
Insert single quotation marks	(As above)	ʹ or ʸ and/or ʹ or ʸ
Insert double quotation marks	(As above)	“ or ” and/or ” or ”
Insert hyphen	(As above)	⊥
Start new paragraph	┌	┌
No new paragraph	┐	┐
Transpose	└┐	└┐
Close up	linking ○ characters	○
Insert or substitute space between characters or words	/ through character or ∧ where required	Υ
Reduce space between characters or words		↑

USING E-ANNOTATION TOOLS FOR ELECTRONIC PROOF CORRECTION

Required Software

Adobe Acrobat Professional or Acrobat Reader (version 7.0 or above) is required to e-annotate PDFs. Acrobat 8 Reader is a free download: <http://www.adobe.com/products/acrobat/readstep2.html>

Once you have Acrobat Reader 8 on your PC and open the proof, you will see the Commenting Toolbar (if it does not appear automatically go to Tools>Commenting>Commenting Toolbar). The Commenting Toolbar looks like this:




If you experience problems annotating files in Adobe Acrobat Reader 9 then you may need to change a preference setting in order to edit.

In the "Documents" category under "Edit – Preferences", please select the category 'Documents' and change the setting "PDF/A mode:" to "Never".



Note Tool — For making notes at specific points in the text

Marks a point on the paper where a note or question needs to be addressed.




How to use it:

1. Right click into area of either inserted text or relevance to note
2. Select Add Note and a yellow speech bubble symbol and text box will appear
3. Type comment into the text box
4. Click the X in the top right hand corner of the note box to close.

Replacement text tool — For deleting one word/section of text and replacing it

Strikes red line through text and opens up a replacement text box.




How to use it:

1. Select cursor from toolbar
2. Highlight word or sentence
3. Right click
4. Select Replace Text (Comment) option
5. Type replacement text in blue box
6. Click outside of the blue box to close

Cross out text tool — For deleting text when there is nothing to replace selection

Strikes through text in a red line.



How to use it:

1. Select cursor from toolbar
2. Highlight word or sentence
3. Right click
4. Select Cross Out Text

Approved tool — For approving a proof and that no corrections at all are required.




How to use it:

1. Click on the Stamp Tool in the toolbar
2. Select the Approved rubber stamp from the 'standard business' selection
3. Click on the text where you want to rubber stamp to appear (usually first page)

Highlight tool — For highlighting selection that should be changed to bold or italic.

Highlights text in yellow and opens up a text box.




How to use it:

1. Select Highlighter Tool from the commenting toolbar
2. Highlight the desired text
3. Add a note detailing the required change

Attach File Tool — For inserting large amounts of text or replacement figures as a files.

Inserts symbol and speech bubble where a file has been inserted.




How to use it:

1. Click on paperclip icon in the commenting toolbar
2. Click where you want to insert the attachment
3. Select the saved file from your PC/network
4. Select appearance of icon (paperclip, graph, attachment or tag) and close

Pencil tool — For circling parts of figures or making freeform marks

Creates freeform shapes with a pencil tool. Particularly with graphics within the proof it may be useful to use the Drawing Markups toolbar. These tools allow you to draw circles, lines and comment on these marks.



How to use it:

1. Select Tools > Drawing Markups > Pencil Tool
2. Draw with the cursor
3. Multiple pieces of pencil annotation can be grouped together
4. Once finished, move the cursor over the shape until an arrowhead appears and right click
5. Select Open Pop-Up Note and type in a details of required change
6. Click the X in the top right hand corner of the note box to close.

Help

For further information on how to annotate proofs click on the Help button to activate a list of instructions:

